

SID 16
(3 hits this paper)

RESULT 2
BTU51820
LOCUS BTU51820 376 bp DNA MAM 16-MAY-1996
DEFINITION Bos taurus Hereford 7 mitochondrial hypervariable region D-loop,
partial sequence.
ACCESSION U51820
VERSION U51820.1 GI:1314616
KEYWORDS .
SOURCE cow strain=Hereford.
ORGANISM Mitochondrion Bos taurus
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea;
Bovidae; Bovinae; Bos.
REFERENCE 1 (bases 1 to 376)
AUTHORS Bradley, D.G., MacHugh, D.E., Cunningham, P. and Loftus, R.T.
TITLE Mitochondrial diversity and the origins of African and European
cattle
JOURNAL Proc. Natl. Acad. Sci. U.S.A. 93 (10), 5131-5135 (1996)
MEDLINE 96209869
REFERENCE 2 (bases 1 to 376)
AUTHORS Bradley, D.G., MacHugh, D.E., Cunningham, P. and Loftus, R.T.
TITLE Direct Submission
JOURNAL Submitted (19-MAR-1996) Daniel G. Bradley, Genetics, Trinity
College Dublin, Dublin, 2, Ireland
FEATURES Location/Qualifiers
source 1. .376
/organism="Bos taurus"
/organelle="mitochondrion"
/strain="Hereford"
/db_xref="taxon:9913"
/note="HE7"
D-loop <1. .>376
/note="hypervariable region"
BASE COUNT 106 a 92 c 60 g 118 t
ORIGIN

Query Match 100.0%; Score 153; DB 3; Length 376;
Best Local Similarity 100.0%; Pred. No. 1e-23;
Matches 153; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 gcatataagcaagtacatgacctctatagcagtacataatacatataattattgactgta 60
|||||
Db 4 GCATATAAGCAAGTACATGACCTCTATAGCAGTACATAATACATATAATTATTGACTGTA 63

Qy 61 catagtacattatgtcaaattcattcttgatagtatatctattatatattccttaccatt 120
|||||
Db 64 CATAGTACATTATGTCAAATTCATTCTTGATAGTATATCTATTATATATTCCTTACCATT 123

Qy 121 agatcacgagcttaattaccatgccgcgtgaaa 153
|||||
Db 124 AGATCACGAGCTTAATTACCATGCCGCGTGAAA 156

sequencer. Consensus sequences were constructed and analyzed with the help of the University of Wisconsin GCG software package. The BLAST algorithm from the National Center for Biotechnology Information (NCBI) was also employed for nucleotide and amino acid sequence homology search (Altschul, S. F. et al. (1990) *J. Mol. Biol.* 215:403-410).

Preparation of Recombinant Proteins:

The recombinant CR8 protein with histidine-tag was prepared using the Xpress™ System (Invitrogen) according to the manufacturer's protocol. cDNA corresponding to the CR8 bHLH domain was obtained by PCR. The sequences of the primers, 5'-GGGGTCTACCAGGGATGTAC-3' (SEQ ID NO:15) for the 5' side, and 5'-GTAAACCACTCTGCAGGGCAATGA-3' (SEQ ID NO:16) for the 3' side, were slightly different from the final consensus sequence for CR8, but the difference did not affect the core bHLH motif. The PCR product was cloned into p17Blue T-vector (Novagen) and subsequently into pRSET-A vector at BamIII and HindIII sites. Constructs were confirmed by DNA sequencing. The protein was overexpressed in JM109 at 37° C. in the presence of isopropylthio- β -D-galactoside (IPTG) by infecting the bacteria with M13 phages that contain the T7 RNA polymerase gene. The cells were lysed with 100 μ g/ml lysozyme in native binding buffer (20 mM sodium phosphate, pH 7.8, 500 mM NaCl), the lysate was loaded on a ProBond™ Ni2+ column, and the recombinant protein was eluted with native-imidazole elution buffer (20 mM sodium phosphate, pH 6.0, 500 mM NaCl, 500 mM imidazole). The protein was then dialyzed against lysis buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA, 10% glycerol, 0.1% Triton X-100, 1 mM DTT) and concentrated on a Microcon 10 (Amicon). To confirm the purity and the quantity, the protein was fractionated on a 12% SDS-polyacrylamide gel with protein standards of known concentrations and visualized by silver staining. The recombinant protein corresponding to the bHLH domain of *da* was supplied by Dr. Michael Caudy (Cornell University Medical College).

Mobility Shift Assay:

The oligonucleotide probes used for the electrophoretic mobility shift assay (EMSA) are as follows: the Class A site used was the CACCTG hexamer (CAGGTG for the opposite strand) from the T5 promoter region of the *Drosophila* AS-C (Villares, R. et al. (1987) *Cell* 50:415-424) (5'-GATCGTAGTCACGCAGGTGGGATCCCTA-3' (SEQ ID NO:17) and 5'-GATCTAGGGATCCACCTGCGTGACTAC-3' (SEQ ID NO:18) for the opposite strand), the Class B site was the CACGTG hexamer from the USF binding site in the adenovirus major late promoter (Gregor, P. D. et al. (1990) *Genes & Dev.* 4:1730-1740) (5'-GATCGGTGTAGGCCACGTGACCGGGTGT-3' (SEQ ID NO:19) and 5'-GATCACACCCGGTCACGTGGCCTACACC-3' (SEQ ID NO:20), the Class C site was the CACGCG hexamer (CGCGTG for the opposite strand) from the AS-C T5 promoter (5'-GATCGGCAGCCGGCACGCGACAGGGCC-3' (SEQ ID NO:21) and 5'-GATCGGCCCTGTGCGGTGCCGGCTGCC-3' (SEQ ID NO:22), and the N-box (CACNAG) was the double hexamer sequence from the Enhancer of split [E(spl)] m8 promoter (Klimbt, C. et al. (1989) *EMBO J.* 8:203-210) (5'-GATCACGCCACGAGCCACAAGGATTG-3' (SEQ ID NO:23) and 5'-GATCCAATCCTTGTGGCTCGTGGCGT-3' (SEQ ID NO:24). One strand of the oligonucleotide was

labeled with [γ -³²P]ATP by T4 polynucleotide kinase, hybridized with three times excess of the opposite strand, and purified using MERmaid oligonucleotide purification kit (BIO 101). 150 ng of the protein was allowed to bind to 50,000 cpm (equivalent to 0.5 ng in a typical experiment) of the labeled probe for 15 min at room temperature in 20 mM HEPES, pH 7.6, 50 mM KCl, 10 mM DTT, 5% glycerol, 0.5 mM EDTA and 0.3 mg/ml BSA. Two microgram of poly (dI-dC) was added to each 20 μ l reaction as on-specific DNA. Samples were analyzed on a 5% native polyacrylamide gel and visualized by autoradiography.

Regulation of CR8:

CR genes were originally defined in IL-2 stimulated normal human T cells. To examine CR8 expression in cytokine-dependent cell lines, the level of CR8 expression was measured by Northern hybridization in the IL-2-dependent human T cell line Kit 225, the IL-3-dependent mouse pro-B cell line Ba/F3, and the IL-2-dependent mouse T cell line CTLL2. The results of this experiment are illustrated in FIGS. 13A-13C. In FIG. 13A, RNA was isolated from quiescent normal human T cells (lanes 1 and 2), IL-2-dependent human T cell line Kit 225 (lanes 3 and 4), IL-3-dependent mouse pro-B cell line Ba/F3 (lanes 5 and 6) and IL-2-dependent mouse T cell line CTLL2 (lanes 7 and 8) left untreated (lanes 1, 3, 5, and 7) or stimulated with 500 pM recombinant human IL-2 (lanes 2, 4, and 8) or 10 U/ml recombinant mouse IL-3 (lane 6) for 2 hr at 37° C. The amount of the growth factor used in the experiment was sufficient to induce maximal [³H]thymidine incorporation. Ten microgram of total RNA was analyzed on formaldehyde/agarose gel and hybridized with either human (lanes 1 to 4) or mouse (lane 5 to 8) CR8 cDNA.

As shown in FIG. 13A, a single 3.2 kb species hybridized to the cDNA probe, and in all three cell lines tested, the level of CR8 was clearly augmented when the cells were stimulated with their respective growth factors. Correlation between the level of CR8 and that of DNA synthesis was in the presence of growth-inhibitory agents was also examined. In this regard, increases in cytoplasmic cAMP are known to inhibit the growth of many cell types, including lymphocytes (Johnson, K. W. et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:6072-6076). IFNs also exert antiproliferative activity on many cell types (Pestka, S. et al. (1987) *Annu. Rev. Biochem.* 56:727-777). Therefore, Kit 225 was stimulated with IL-2, IFN β , or forskolin, which increases cytoplasmic cAMP by activating adenylate cyclase, either alone or in combination. IL-2-dependent [³H]thymidine incorporation was inhibited by IFN β and forskolin in Kit 225 cells in a dose-dependent fashion (FIG. 13B). FIG. 13B demonstrates that IFN β and forskolin inhibit IL-2-dependent [³H]thymidine incorporation by Kit 225 cells. Ten thousand quiescent Kit 225 cells were incubated with indicated reagents in 200 μ l for 24 hr at 37° C. The culture was pulsed with [³H]thymidine for the last 4 hr to monitor the DNA synthesis. (●), IL-2 only (500 pM); (◇), IL-2 500 pM+varying concentrations of IFN β (U/ml); (○), IL-2 500 pM+varying concentrations of forskolin (fM). While forskolin was capable of reducing the IL-2-dependent [³H]thymidine incorporation almost to the basal level, IFN β -mediated inhibition never exceeded 70% of the maximal incorporation in several independent experiments. The expression of CR8 was compared with that of c-myc, an IL-2-inducible immediate-early gene that encodes a bHLH protein and is implicated for cell proliferation (Marcu, K. B. et al. (1992) *Annu. Rev. Biochem.* 61:809-860).

FIG. 13C shows the effect of antiproliferative agents on the expression of CR8 and c-myc transcripts in Kit 225.

5104

RESULT 7
 BTU51824
 LOCUS BTU51824 389 bp DNA MAM 16-MAY-1996
 DEFINITION Bos taurus Jersey 5 mitochondrial hypervariable region D-loop, partial sequence.
 ACCESSION U51824
 VERSION U51824.1 GI:1314620
 KEYWORDS .
 SOURCE cow strain=Jersey.
 ORGANISM Mitochondrion Bos taurus
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Cetartiodactyla; Ruminantia; Pecora; Bovoidea; Bovidae; Bovinae; Bos.
 REFERENCE 1 (bases 1 to 389)
 AUTHORS Bradley,D.G., MacHugh,D.E., Cunningham,P. and Loftus,R.T.
 TITLE Mitochondrial diversity and the origins of African and European cattle
 JOURNAL Proc. Natl. Acad. Sci. U.S.A. 93 (10), 5131-5135 (1996)
 MEDLINE 96209869
 REFERENCE 2 (bases 1 to 389)
 AUTHORS Bradley,D.G., MacHugh,D.E., Cunningham,P. and Loftus,R.T.
 TITLE Direct Submission
 JOURNAL Submitted (19-MAR-1996) Daniel G. Bradley, Genetics, Trinity College Dublin, Dublin, 2, Ireland
 FEATURES Location/Qualifiers
 source 1. .389
 /organism="Bos taurus"
 /organelle="mitochondrion"
 /strain="Jersey"
 /db_xref="taxon:9913"
 /note="JE5"
 D-loop <1. .>389
 /note="hypervariable region"
 BASE COUNT 111 a 93 c 61 g 124 t
 ORIGIN

Query Match 100.0%; Score 46; DB 3; Length 389;
 Best Local Similarity 100.0%; Pred. No. 1.8e-08;
 Matches 46; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy 1 attatatgccccatgcatataagcaagtacatgacctctatagcag 46
 |||||
 Db 3 ATTATATGCCCCATGCATATAAGCAAGTACATGACCTCTATAGCAG 48

2 111 from 1996 paper

-continued

AAGCCAGCCG GGAAGAGAGG TTGAGTCACC CATCCCCGAG GTGGCTGCCC CTGTGTGGGA 002
 GGCAGGTTCT GCAAAGCAAG TGCAAGAGGA CAAAAA AAAA AAAAATGCGC 862
 TCCAGCAGCC TGTTTGGGAA GCAGCAGTCT CTCCTTCAGA TACTGTGGGA CTCATGCTGG 922
 AGAGGAGCCG CCCACTTCCA GGACCTGTGA ATAAGGGCTA ATGATGAGGG TTGGTGGGGC 982
 TCTCTGTGGG GCAAAAAGGT GGTATGGGGG TTAGCACTGG CTCTCGTTCT CACCGAGAGAA 1042
 GGAAGTGTTC TAGTGTGGTT TAGGAAACAT GTGGATAAAG GGAACCATGA AAATGAGAGG 1102
 AGGAAAGACA TCCAGATCAG CTGTTTGGCC TGTGCTCAG TTGACTCTGA TTGCATCCTG 1162
 TTTTCCTAAT TCCCAGACTG TTCTGGGCAC GGAAGGGACC CTGGATGTGG AGTCTTCCCC 1222
 TTTGGCCCTC CTCAGTGGCC TCTGGGCTAG CCCAGAGTCC CTTAGCTTGT ACCTCGTAAC 1282
 ACTCCTGTGT GTCTGTCCAG CCTTGCAGTC ATGTCAAGGC CAGCAAGCTG ATGTGACTCT 1342
 TGCCCCATGC GAGATATTTA TACCTCAAAC ACTGGCCTGT GAGCCCTTTC CAACTCAGTG 1402
 GAGAGCCCTG AAAGGAGCCT CACTTGAATC CAGCTCAGTG CTCTGGGTGG CCCCTGCAG 1462
 GTGCCCCCTG ACCCTGCGTT GCAGCAGGGT CCACCTGTGA GCAGGCCCGC CCTGGGCCCT 1522
 CTTCTGGAT GTGCCCTCTC TGAGTTCTGT GCTGTCTCTT GGAGGCAGGG CCCAGGAGAA 1582
 CAAAGTGTGG AGGCCTCGGG GAGTGACTTT TCCAGCTCTC ATGCCCCGCA GTGTGAACA 1642
 AGGCAGAAAA GGATCCTAGG AAATAAGTCT CTTGGCGGTC CCTCAGAGTC CTGCTGAAAT 1702
 CCAGCCAGTG TTTTTTGTGG TATGAGAACA GCCAAAAGA GATGCCCGA GATAGAAGGG 1762
 GAGCCTTG TGTTCTTCTC GCAGACGTGA GATGAACACT GGAGTGGGCA GAGGTGGCCC 1822
 AGGACCATGA CACCCTTAGA GTGCAGAAGC TGGGGGAGA GGCTGCTTCG AAGGGCAGGA 1882
 CTGGGGATAA TCAGAACCTG CCTGTACCT CAGGGCATCA CTGAACAAAC ATTTCTGAT 1942
 GGGAACTCT GCGGCAGAGC CCAGGCTGGG GAAGTGAAC ACCCAGGCA GCCCCTTTGT 2002
 GGCCAGGAT AATCAACACT GTTCTCTCTG TACCATGAGC TCCTCCAGGA GATTATTAA 2062
 GTGTATTGTA TCATTGGTTT TCTGTGATTG TCATAACATT GTTTTGTTA CTGTTGGTGC 2122
 TGTGTATT TATTATTGTA ATTTCAGTTT GCCTTACTG GAGAATCTCA GCAGGGGTTT 2182
 CAGCCTGACT GTCTCCCTTT CTCTACCAGA CTCTACCTCT GAATGTGCTG GGAACCTCTT 2242
 GGAGCCTGTC AGGAACTCCT CACTGTTTAA ATATTAGGT ATTGTGACAA ATGGAGCTGG 2302
 TTTCTAGAA ATGAATGATG TTTGCAATCC CCATTTTCCT GTTTCAGCAT GTTATATTCT 2362
 TATGAAATAA AAGCCCAAGT CCAATATGAA AAAAAAAAAA AAAA 2406

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 202 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Cys Arg Thr Leu Ala Ala Phe Pro Thr Cys Leu Glu Arg Ala
 1 5 10 15
 Lys Glu Phe Lys Thr Arg Leu Gly Ile Phe Leu His Lys Ser Glu Leu
 20 25 30
 Gly Cys Asp Thr Gly Ser Thr Gly Lys Phe Glu Trp Gly Ser Lys His
 35 40 45
 Ser Lys Glu Asn Arg Asn Phe Ser Glu Asp Val Leu Gly Trp Arg Glu
 50 55 60